

FORM TP 2008241



TEST CODE **22134032**

MAY/JUNE 2008

Trinidad & Tobago

CARIBBEAN EXAMINATIONS COUNCIL
ADVANCED PROFICIENCY EXAMINATION

PURE MATHEMATICS

UNIT 1 – PAPER 03/B

ALGEBRA, GEOMETRY AND CALCULUS

1 ½ hours

26 JUNE 2008 (a.m.)

This examination paper consists of **THREE** sections: Module 1, Module 2 and Module 3.

Each section consists of 1 question.

The maximum mark for each Module is 20.

The maximum mark for this examination is 60.

This examination consists of 4 printed pages.

INSTRUCTIONS TO CANDIDATES

1. **DO NOT** open this examination paper until instructed to do so.
2. Answer **ALL** questions from the **THREE** sections.
3. Write your solutions, with full working, in the answer booklet provided.
4. Unless otherwise stated in the question, any numerical answer that is not exact **MUST** be written correct to three significant figures.

Examination Materials Permitted

Graph paper (provided)

Mathematical formulae and tables (provided) – **Revised 2008**

Mathematical instruments

Silent, non-programmable, electronic calculator


Copyright © 2008 Caribbean Examinations Council ®.
All rights reserved.

SECTION A (Module 1)

Answer this question.

1. (a) (i) Write $\log_2 2^p$ in terms of p only. **[2 marks]**
- (ii) Solve for x the equation $\log_2 [\log_2 (2x - 2)] = 2$. **[3 marks]**
- (b) The diagram below (**not drawn to scale**) shows the graph of the function

$$f(x) = 3x^3 + hx^2 + kx + m \text{ which touches the } x\text{-axis at } x = -1.$$



- (i) Determine the values of the constants h , k and m . **[7 marks]**
- (ii) State the range of values of x in $(-\infty, 0]$ for which $f(x)$ is a decreasing function. **[2 marks]**
- (c) Evaluate $\sum_{r=1}^{100} (3r + 2)$. **[6 marks]**


Total 20 marks

GO ON TO THE NEXT PAGE

SECTION B (Module 2)

Answer this question.

2. The diagram below shows the path of a comet around the sun S. The path is described by the parametric equation $x = at^2$ and $y = 2at$, where $a > 0$ is a constant.



- (a) Show that the Cartesian equation for the path is $y^2 = 4ax$. **[2 marks]**
- (b) Given that the gradient m of the tangent at any point on the path satisfies $m = \frac{2a}{y}$,
- (i) show that the equation of the tangent at (x_1, y_1) is $yy_1 = 2a(x + x_1)$ in Cartesian form and $ty = x + at^2$ in parametric form **[5 marks]**
- (ii) find the equation of the normal at the point P with parameter t_1 **[3 marks]**
- (iii) show that $t_1^2 + t_1 t_2 + 2 = 0$ if the normal in (ii) above intersects the path again at the point P' with parameter t_2 **[6 marks]**
- (iv) find the distance $|QR|$ if the tangent at P meets the x -axis at Q and the normal meets the x -axis at R. **[4 marks]**

Total 20 marks

GO ON TO THE NEXT PAGE

SECTION C (Module 3)

Answer this question.

3. (a) (i) By expressing $x - 9$ as $(\sqrt{x} + 3)(\sqrt{x} - 3)$, find $\lim_{x \rightarrow 9} \frac{\sqrt{x} - 3}{x - 9}$. [3 marks]


(ii) Hence, find $\lim_{x \rightarrow 9} \frac{\sqrt{x} - 3}{x^2 - 10x + 9}$. [4 marks]

(b) (i) Find the value of u if $\int_u^{2u} \frac{1}{x^2} dx = \frac{1}{4}$. [3 marks]

(ii) Given that $\int_1^4 f(x) dx = 7$, evaluate

$$\int_1^2 [f(x) + 1] dx + \int_2^4 [f(x) - 2] dx. \quad [5 \text{ marks}]$$

(c) The figure below (not drawn to scale) shows a hemispherical bowl which contains liquid.



The volume $V \text{ cm}^3$ of liquid is given by

$$V = \frac{1}{3} \pi h^2 (24 - h)$$

where h is the greatest depth of the liquid in cm. Liquid is poured into the bowl at the rate of 100 cm^3 per second.

(i) Find $\frac{dV}{dt}$ in terms of h . [3 marks]

(ii) Calculate the rate at which h is increasing when $h = 2$ cm. (Leave your answer in terms of π .) [2 marks]

Total 20 marks



END OF TEST